



This Product Information Sheet is not intended to replace the Product Label. Always read the Product Label for the most current nutrient content and feeding directions.

Breed-Up® 17 Pro Mag with Garlic and ClariFly®

Product #83656

Feeding Applications

For beef cattle on pasture

Ingredients

Molasses Products, Plant Protein Products, Magnesium Oxide, Monocalcium Phosphate, Dicalcium Phosphate, Processed Grain By-Products, Vegetable Oil, Calcium Carbonate, Roughage Products, Manganese Sulfate, Zinc Proteinate, Manganese Proteinate, Mineral Oil, Zinc Sulfate, Selenium Yeast, Copper Sulfate, Silicon Dioxide, Natural Flavor, Manganous Oxide, Copper Proteinate, Cobalt Proteinate, Zinc Oxide, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Ethylenediamine Dihydroiodide, Cobalt Carbonate, Calcium Iodate.

Nutrition Information

Crude Protein (min)	17%
Crude Fat (min)	3%
Crude Fiber (max)	3%
Calcium	1% - 1.5%
Phosphorus (min)	1%
Salt (min)	None Added
Magnesium (min)	4%
Potassium (min)	1.5%
Vitamin A (min)	100000 IU/lb
Vitamin D3 (min)	10000 IU/lb
Vitamin E (min)	100 IU/lb
Cobalt (min)	13 ppm
Copper (min)	400 ppm
Iodine (min)	30 ppm
Manganese (min)	1350 ppm
Selenium (min)	6 ppm
Zinc (min)	1200 ppm

Usage Guidelines

Placement: Locate where cattle congregate (watering, loafing, shade areas).

Quantity: Offer cattle one block for every 15 to 25 head. Cattle should consume an average of 0.8 oz per 100 lbs of animal body weight per day, equivalent to 8 oz per day for a 1,000-lb cow.

Additional Feeding Information

Prevents the emergence of house flies, horn flies, face flies and stable flies from manure of treated cattle.

Active Ingredient: Diflubenzuron (CAS #35367-38-5) 0.02% (contains 90.8 mg of diflubenzuron per pound). Diflubenzuron Group 15 Insecticide. EPA Reg. No. 89459-7-7698. EPA Est. No. 7698-KY-002H, 7698-MN-003B, 7698-SD-003A, 7698-TX-001D.

Provide as a free choice source of minerals for your cattle. Start feeding early in the spring before flies appear and continue through summer and into fall until cold weather restricts fly activity. Existing adult flies will not be affected. Ensure cattle are not starved for nutrients prior to feeding. Remove all other self-limiting nutritional supplements. Do not use in association with feeds containing supplemental selenium. If intake is below 0.8 oz per 100 lb body weight per day, increase or relocate feeding locations to areas more frequented by cattle; if above, reduce or relocate to less-frequented areas. Introduction after adult fly infestation is established may require supplemental sprays.

WARNING: This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.