This Product Information Sheet is not intended to replace the Product Label. Always read the Product Label for the most current nutrient content and feeding directions.

Crystalyx 25% Beef

Product #3799

Feeding Applications

For all classes of beef cattle

Product Features

CRYSTALYX® 25% Beef combines both rumen degradable protein and rumen bypass protein. This nutrient dense self-fed supplement provides protein, minerals and vitamins for cattle on low to moderate quality forages.

- Consistent consumption, even in challenging conditions
- Formulated with Bioplex Organic Trace Minerals: A chelated form of trace minerals that are more readily absorbed and available to be used by the animal
- 2% Magnesium oxide for added magnesium supplementation
- Phosphorus at 1% will supply adequate phosphorus on most forages through the first trimester of gestation for a beef cow. After that, additional mineral (phosphorus) supplementation is recommended through peak lactation or breeding.

Product availability: Available only in CA, OR, WA and NV.

Ingredients

Molasses Products, Processed Grain By-Products, Urea, Calcium Phosphate, Magnesium Oxide, Calcium Carbonate, Vegetable Oil, Plant Protein Products, Zinc Proteinate, Manganese Sulfate, Manganese Proteinate, Copper Sulfate, Selenium Yeast, Roughage Products, Zinc Sulfate, Copper Proteinate, Manganous Oxide, Cobalt Proteinate, Zinc Oxide, Vitamin E Supplement,, Vitamin A Supplement, Vitamin D3 Supplement, Ethylenediamine Dihydroiodide, Mineral Oil, Cobalt Carbonate, Calcium Iodate.

Nutrition Information

Crude Protein (min)	25%
Crude Fat (min)	4%
Crude Fiber (max)	2.5%
Calcium	1.5% - 2%
Phosphorus (min)	1%
Magnesium (min)	2%
Potassium (min)	2%
Vitamin A (min)	50000 IU/lb
Vitamin D3 (min)	5000 IU/lb
Vitamin E (min)	50 IU/lb
Cobalt (min)	15 ppm
Copper (min)	500 ppm
Iodine (min)	17 ppm
Manganese (min)	1330 ppm
Selenium (min)	6 ppm
Zinc (min)	1000 ppm

Usage Guidelines

Placement: Place containers in areas that achieve desired supplement intake.

Relocation: Once acceptable intake levels have been achieved, the containers may be moved to underutilized pasture areas to improve grazing distribution and forage utilization.

Accessibility: Feed at the rate of one container for every 20-30 head. A minimum of two (2) containers may be necessary in each free-stall pen, lot or pasture to ensure proper accessibility.

Monitoring: Self-fed consumption will depend on size of the animals fed, seasonal weather conditions, as well as quality, source and availability of other feeds. Cattle typically consume between 0.5 and 1.1 pounds per head per day.